

2022 Black Shank Fungicide Trial for Dark Tobacco

Kent Boyd Farm – Hopkinsville KY


- KTD6 transplanted June 3
 - Moderate black shank resistance (3-3).
- Fungicide applications:
 - Transplant Water (TPW, 200 gal/A)
 - 3 weeks after transplanting
 - Banded spray at 15 gal/A
 - 5 weeks after transplanting
 - Banded spray at 15 gal/A
- Tobacco harvested late September, fire-cured.
- Treatments:
 - 1) Ridomil Gold (8 oz/A) TPW
 - 2) Orondis Gold (4.8 oz/A) + Ridomil Gold (8 oz/A) TPW
 - 3) Presidio (4 oz/A) at 3 wks
 - 4) Presidio (4 oz/A) at 3 wks, Ridomil Gold (16 oz/A) at 5 wks
 - 5) Orondis Gold (4.8 oz/A) + Ridomil Gold (8 oz/A) TPW, Presidio (4 oz/A) at 3 wks, Ridomil Gold (16 oz/A) at 5 wks
 - 6) Untreated Control

2022 Black Shank Fungicide Trial for Dark Tobacco

Kent Boyd Farm – Hopkinsville KY

LSD_{0.10} = 31.1

■ Final Stand (% survival)



2022 Black Shank Fungicide Trial for Dark Tobacco

Kent Boyd Farm – Hopkinsville KY

LSD_{0.10} =

189 807
■ Lug ■ Leaf

980 (total)

